

PISD Athletic Department

Guidelines for Concussion Management

Introduction

Approximately 10 percent of all athletes involved in contact sports suffer a Mild Traumatic Brain Injury (concussion) each season; some estimates are as high as 19 percent. Because many mild concussions can go undiagnosed and unreported, it is difficult to estimate precisely the rate of concussion in any sport. Symptoms are not always definite, and knowing when it is safe for an athlete to return to play is not always clear.

The recognition and management of concussion in athletes can be difficult for a number of reasons:

Athletes who have experienced a concussion can display a wide variety of symptoms. Although the classic symptoms of loss of consciousness, confusion, memory loss, and/or balance problems may be present in some athletes with mild traumatic brain injury, there may or may not be obvious signs that a concussion has occurred.

Post-concussion symptoms can be quite subtle and may go unnoticed by the athlete, team medical staff, or coaches. Many coaches and other team personnel may have limited training in recognizing signs of concussion and therefore may not accurately diagnose the injury when it has occurred. Players may be reluctant to report concussive symptoms for fear that they will be removed from the game, and this may jeopardize their status on the team, or their athletic careers.

Pearland ISD is in compliance with HB 2038, 82(R).

- A student removed from an athletics practice or competition would not be permitted to practice or compete again until the student has been evaluated and cleared to play through a school-issued written statement by the treating physician.
- The student's parent or guardian and student would have to return the physician's statement and complete a consent form indicating that they had been informed and consented to the policies established under the return-to-play protocol.
- Parents and student understands the risks associated with the student's returning to play and would comply with any ongoing requirements outlined by the concussion policy.
- Parents must consent to the physician's disclosure of health information that is related to the concussion treatments.
- And, parents understand the district or school's immunity from liability provisions.

The Pearland ISD Concussion Oversight Team includes:

Chris Shaddock, LAT, ATC- Athletic Trainer
Savannah Tracy, LAT, ATC- Athletic Trainer
Christina Decoteau, LAT, ATC- Athletic Trainer
Matt Thomas, MS, LAT, ATC- Athletic Trainer
William Ryan, MAT, LAT, ATC- Athletic Trainer
Jill Flowers, LAT, ATC- Athletic Trainer
Jason Decker, MD – Team Physician
Taggart T. Gauvain, MD Team Physician
Evan Meeks, MD – Team Physician
Summer Ott, Psy.D. – Neuropsychologist

Recovery and safe return-to-play

It is crucial to allow enough healing and recovery time following a concussion to prevent further damage. Research suggests that the effects of repeated concussion are cumulative over time.

Most athletes who experience an initial concussion can recover completely as long as they do not return to contact sports too soon. Following a concussion, there is a period of change in brain function that may last anywhere from 24 hours to 10 days or longer. During this time, the brain may be vulnerable to more severe or permanent injury. If the athlete sustains a second concussion during this time period, the risk of permanent brain injury increases.

Definitions

Concussion or Mild Traumatic Brain Injury (MTBI) - A concussion or MTBI is the common result of a blow to the head or body which causes the brain to move rapidly within the skull. This injury causes brain function to change which results in an altered mental state (either temporary or prolonged). Physiologic and/or anatomic disruptions of connections between some nerve cells in the brain occur. Concussions can have serious and long-term health effects, even from a mild bump on the head. Symptoms include, but are not limited to, headache, amnesia, nausea, dizziness, confusion, blurred vision, ringing in the ears, loss of balance, moodiness, poor concentration or mentally slow, lethargy, photosensitivity, sensitivity to noise, and a change in sleeping patterns. Symptoms can also include a loss of consciousness but many do not. These symptoms may be temporary or long lasting.

Second Impact Syndrome – Second impact syndrome (SIS) refers to catastrophic events which may occur when a second concussion occurs while the athlete is still symptomatic and healing from a previous concussion. The second injury may occur within days or weeks following the first injury. Loss of consciousness is not required. The second impact is more likely to cause brain swelling with other widespread damage to the brain. This can be fatal. Most often SIS occurs when an athlete returns to activity without being symptom free from the previous concussion.

Prevention Strategies

Helmets, headgear, and mouth guards do not prevent all concussions.

1. All headgear must be NOCSAE certified.
2. Make sure the headgear fits the individual.
3. For all sports that require headgear, a coach or appropriate designate should check headgear before use to make sure air bladders work and are appropriately filled. Padding should be checked to make sure they are in proper working condition.
4. Make sure helmets are secured properly at all times.
5. Mouth guards should fit and be used at all times.

Evaluation for Concussion/MTBI

1. At time of injury administer one of these assessment tests:
 - a. Sports Concussion Assessment Tool (SCAT 5)
 - b. Graded Symptom Checklist (GSC)
2. Observe athlete 15 to 20 minutes and re-evaluate.
3. Athlete does not return to a game or practice if he/she has any signs or symptoms of Mild Traumatic Brain Injury (Concussion).
4. Doctor Referral
5. Home Instructions
6. Return to Play Guidelines for Parents
7. **Note - If in doubt, athlete is referred to physician and does not return to play.**

Concussion Management

1. Recommended school modifications
 - a. Notify Nurse, Assistant Principal and Counselor of the student that he/she has MTBI
 - b. Notify Nurse, Assistant Principal and Counselor of post-concussion symptoms
 - c. Student may need special accommodations such as limited computer work, reading activities, testing, assistance to class, etc. until symptoms subside
 - d. Student may only be able to attend school for half days or may need daily rest periods until symptoms subside with physician authorization
2. The treating physician must provide a written statement to the parent and athletic trainer indicating that, in the physician's Professional judgment, it is safe for the student to return to play.
3. Student must show no signs of post-concussion symptoms before return to play protocol begins, unless under physician guidelines for a submaximal exercise program.
4. Student will not return to full practice or competition for minimum of 7 days.
5. Student athlete and the parent/guardian have signed the form acknowledging the completion of the return to play guidelines which includes the understanding the risks associated with the student athlete's return to play.

Return to Play Guidelines

Athlete must stay in Phase I (Rest/Physician Clearance) for a minimum of 24-48 hours before continuing return to play protocol or beginning a submaximal program.

1. Athlete activity progressions (There must be 24 hours of rest between progression phases)
 - I. Rest / Physician clearance
 - II. Light aerobic exercise with no resistance training *or physician prescribed submaximal program*
 - III. Moderate aerobic activity with resistance training
 - IV. Sport specific activity and Non-contact training drills
 - V. Full contact training drills can begin after minimum 7 days
 - VI. Return to full participation (pending physician clearance)

Note – Athlete activity progression continues as long as athlete is asymptomatic at current level. If the athlete experiences any post-concussion symptoms, stop physical activity until symptom free for 24-48 hours. Resume with phase or level in which they were previously asymptomatic. If an athlete is operating under a prescribed submaximal program, the athlete must follow doctor's orders for symptomatic exercise. An athlete may not participate in submaximal program while experiencing balance problems, dizziness, or vision problems.

2. Physician clearance
3. Athletic Trainer clearance

Home Instructions

Following these home instructions can prevent further injury and help in recovery.

WHEN TO SEEK CARE URGENTLY:

Seek care quickly if symptoms worsen and watch for any of the following danger signs. If you observe any of the following, call your doctor or proceed to the nearest emergency department:

Headaches that worsen Seizures Neck Pain Unusual behavior change	Very drowsy, can't be awakened Repeated vomiting Slurred speech Significant irritability	Can't recognize people or places Increasing confusion Weakness/numbness in arms/legs Less responsive than usual
---	---	--

COMMON SIGNS & SYMPTOMS

With a concussion, it is common to have one or many symptoms including physical, cognitive, emotional, and sleep irregularities. Record all symptoms on the graded symptom checklist.

Physical		Cognitive	Emotional	Sleep
Headache	Visual Problems	Feeling mentally foggy	Irritability	Drowsiness
Nausea/Vomiting	Fatigue	Feeling slowed down	Sadness	Sleeping less than usual
Dizziness	Sensitive to light or noise	Difficulty remembering	More Emotional	Sleeping more than usual
Balance Problems	Numbness/Tingling	Difficulty concentrating	Nervousness	Trouble falling asleep

RETURNING TO DAILY ACTIVITIES

The key to recovery is sleeping, resting physically and mentally, and avoiding activities that might cause another head injury.

- **Avoid/Limit:**
 - Physical activities such as PE, sport practice, weight training, running, exercising, heavy lifting, etc.
 - Lengthy mental activities requiring concentration (i.e. reading, schoolwork, job-related work, phone use, and video games) as these activities worsen symptoms and prolong recovery
- **Sleep:**
 - Get a lot of rest. Be sure to get enough sleep at night – no late nights. Keep the same bedtime weekdays & weekends
 - Take daytime naps or rest breaks when you feel tired or fatigued
 - It is not necessary to wake up periodically
- **Nutrition:**
 - Drink lots of fluids and eat a well-balanced diet
- **Evaluate:**
 - Report your symptoms daily to help guide recovery

DO'S & DON'TS

It is OK to:	There is NO need to:	Do Not
Take acetaminophen (Tylenol) Use ice packs for head and neck Go to Sleep Rest	Stay in Bed Wake every hour	Drive with symptoms Exercise or lift weights Drink Alcohol Participate in sports

DHS: 281-412-8835

Chris Shaddock, LAT, ATC
shaddockc@pearlandisd.org
Christina Decoteau, MS, LAT, ATC
decoteauc@pearlandisd.org
Savannah Tracy, LAT, ATC
tracys@pearlandisd.org

PHS: 281-997-3260

Matt Thomas, MS, LAT, ATC
thomasma@pearlandisd.org
Jill Flowers, LAT, ATC
flowersj@pearlandisd.org
Will Ryan, MAT, LAT, ATC
ryanw@pearlandisd.org

BMJH & PJHS

Brittnie Smithley, MAT, LAT, ATC
smithleyb@pearlandisd.org

PJHE & PJHW

Kathy Nguyen, MAT, LAT, ATC
nguyenku@pearlandisd.org

Parent Name
(Please Print)

Parent Signature

Date

PEARLAND ISD HEAD INJURY RETURN TO PLAY FORM

Name of Student: _____ Sport: _____ School: _____ Date of Injury: _____

This form must be completed and submitted to the athletic trainer or other person (who is not a coach) responsible for compliance with the Return to Play protocol established by the school district Concussion Oversight Team, as determined by the superintendent or their designee (see Section 38.157 (c) of the Texas Education Code).

Parent/Guardian signs and certifies that he/she:

Has been informed concerning and consents to the student participating in returning to play in accordance with the return to play protocol established by the Concussion Oversight Team. Understands the risks associated with the student returning to play and will comply with any ongoing requirements in the return to play protocol. Consents to the disclosure to appropriate persons, consistent with the Health Insurance Portability and Accountability Act of 1996 (Pub. L. No. 104-191), of the treating physician's written statement under Subdivision (3) and, if any, the return to play recommendations of the treating physician. Understands the immunity provisions under Section 38.159 of the Texas Education Code.

Parent/Guardian Printed Name

Parent/Guardian Signature

Date

Athletic Trainer verifies:

The student has been evaluated by a treating physician selected by the student, their parent or other person with legal authority to make medical decisions for the student. The student has completed the Return to Play protocol established by the school district Concussion Oversight Team. The school has received a written statement from the treating physician indicating, that in the physician's professional judgment, it is safe for the student to return to play.

Athletic Trainer Printed Name

Athletic Trainer Signature

Date

RETURN TO PLAY GUIDELINES

Athletes must complete the following stepwise process prior to return to play following a concussion: No activity and rest until symptom free and there must be 24 hours between each phase, unless under a prescribed submaximal program.

PHASE 1 - Rest

PHASE 2 - Light aerobic exercise *or physician prescribed submaximal program*

PHASE 3 - Moderate aerobic exercise with resistance training

PHASE 4 - Sport-specific training / Noncontact drills

PHASE 5 - Full-contact drills (minimum 7 days

post-injury) PHASE 6 - Game play

Note – Athlete activity progression continues as long as athlete is asymptomatic at current level. If the athlete experiences any post-concussion symptoms, stop physical activity until symptom free for 24-48 hours. Resume with phase or level in which they were previously asymptomatic. If an athlete is operating under a prescribed submaximal program, the athlete must follow doctor's orders for symptomatic exercise. An athlete may not participate in submaximal program while experiencing balance problems, dizziness, or vision problems.

PHYSICIAN RECOMMENDATIONS

Returning to School

- No restrictions
- Full Days as tolerated
- Half days as tolerated until _____
- No school
 - May return on ____/____/____ for FULL days
 - May return on ____/____/____ for HALF days
- Homebound instruction _____ hours per week
- Postpone exams
- No more than __ exams/week
- Reduced workload (i.e. assign ½ homework problems)
- Allow frequent breaks. May require quiet area separate from others
- Make up missed assignments gradually
- Provide preprinted class notes / allow to obtain notes from peers
- Tutoring as needed

Signs observed at time of injury

- [] Appears to be dazed or stunned
- [] Is confused about assignment
- [] Forgets plays
- [] Is unsure of game, score, or opponent
- [] Moves clumsily
- [] Answers questions slowly
- [] Loss of consciousness (even temporarily)
- [] Shows behavior or personality change
- [] Forgets events prior to hit (retrograde amnesia)
- [] Forgets events after hit (anterograde amnesia)

Symptoms reported by athlete at time of injury

- [] Headache
- [] Nausea
- [] Balance problems or dizziness
- [] Double or fuzzy vision
- [] Sensitivity to light or noise
- [] Feeling sluggish
- [] Feeling "foggy"
- [] Change in sleep pattern
- [] Concentration or memory problems

Returning to Activity

- __ Begin submaximal exercise as indicated on the attached guidelines
- __ Begin Return to Play Protocol when symptom free
- __ Begin Return to Play Protocol on ____/____/____
- __ Begin Return to Play Protocol after ____ days symptom free
- __ No activity until follow up on ____/____/____

Physician's Signature: _____ **Date:** _____

Physician's Address: _____ **Phone:** _____

SCAT5[®]

SPORT CONCUSSION ASSESSMENT TOOL – 5TH EDITION

DEVELOPED BY THE CONCUSSION IN SPORT GROUP

FOR USE BY MEDICAL PROFESSIONALS ONLY

supported by



Patient details

Name: _____

DOB: _____

Address: _____

ID number: _____

Examiner: _____

Date of Injury: _____ Time: _____

WHAT IS THE SCAT5?

The SCAT5 is a standardized tool for evaluating concussions designed for use by physicians and licensed healthcare professionals¹. The SCAT5 cannot be performed correctly in less than 10 minutes.

If you are not a physician or licensed healthcare professional, please use the Concussion Recognition Tool 5 (CRT5). The SCAT5 is to be used for evaluating athletes aged 13 years and older. For children aged 12 years or younger, please use the Child SCAT5.

Preseason SCAT5 baseline testing can be useful for interpreting post-injury test scores, but is not required for that purpose. Detailed instructions for use of the SCAT5 are provided on page 7. Please read through these instructions carefully before testing the athlete. Brief verbal instructions for each test are given in italics. The only equipment required for the tester is a watch or timer.

This tool may be freely copied in its current form for distribution to individuals, teams, groups and organizations. It should not be altered in any way, re-branded or sold for commercial gain. Any revision, translation or reproduction in a digital form requires specific approval by the Concussion in Sport Group.

Recognise and Remove

A head impact by either a direct blow or indirect transmission of force can be associated with a serious and potentially fatal brain injury. If there are significant concerns, including any of the red flags listed in Box 1, then activation of emergency procedures and urgent transport to the nearest hospital should be arranged.

Key points

- Any athlete with suspected concussion should be **REMOVED FROM PLAY**, medically assessed and monitored for deterioration. No athlete diagnosed with concussion should be returned to play on the day of injury.
- If an athlete is suspected of having a concussion and medical personnel are not immediately available, the athlete should be referred to a medical facility for urgent assessment.
- Athletes with suspected concussion should not drink alcohol, use recreational drugs and should not drive a motor vehicle until cleared to do so by a medical professional.
- Concussion signs and symptoms evolve over time and it is important to consider repeat evaluation in the assessment of concussion.
- The diagnosis of a concussion is a clinical judgment, made by a medical professional. The SCAT5 should **NOT** be used by itself to make, or exclude, the diagnosis of concussion. An athlete may have a concussion even if their SCAT5 is "normal".

Remember:

- The basic principles of first aid (danger, response, airway, breathing, circulation) should be followed.
- Do not attempt to move the athlete (other than that required for airway management) unless trained to do so.
- Assessment for a spinal cord injury is a critical part of the initial on-field assessment.
- Do not remove a helmet or any other equipment unless trained to do so safely.

IMMEDIATE OR ON-FIELD ASSESSMENT

The following elements should be assessed for all athletes who are suspected of having a concussion prior to proceeding to the neurocognitive assessment and ideally should be done on-field after the first first aid / emergency care priorities are completed.

If any of the "Red Flags" or observable signs are noted after a direct or indirect blow to the head, the athlete should be immediately and safely removed from participation and evaluated by a physician or licensed healthcare professional.

Consideration of transportation to a medical facility should be at the discretion of the physician or licensed healthcare professional.

The GCS is important as a standard measure for all patients and can be done serially if necessary in the event of deterioration in conscious state. The Maddocks questions and cervical spine exam are critical steps of the immediate assessment; however, these do not need to be done serially.

STEP 1: RED FLAGS

RED FLAGS:

- Neck pain or tenderness
- Double vision
- Weakness or tingling/burning in arms or legs
- Severe or increasing headache
- Seizure or convulsion
- Loss of consciousness
- Deteriorating conscious state
- Vomiting
- Increasingly restless, agitated or combative

STEP 2: OBSERVABLE SIGNS

Witnessed Observed on Video

	Y	N
Lying motionless on the playing surface	Y	N
Balance / gait difficulties / motor incoordination: stumbling, slow / laboured movements	Y	N
Disorientation or confusion, or an inability to respond appropriately to questions	Y	N
Blank or vacant look	Y	N
Facial injury after head trauma	Y	N

STEP 3: MEMORY ASSESSMENT MADDOCKS QUESTIONS²

"I am going to ask you a few questions, please listen carefully and give your best effort. First, tell me what happened?"

Mark Y for correct answer / N for incorrect

	Y	N
What venue are we at today?	Y	N
Which half is it now?	Y	N
Who scored last in this match?	Y	N
What team did you play last week / game?	Y	N
Did your team win the last game?	Y	N

Note: Appropriate sport-specific questions may be substituted.

Name: _____
 DOB: _____
 Address: _____
 ID number: _____
 Examiner: _____
 Date: _____

STEP 4: EXAMINATION

GLASGOW COMA SCALE (GCS)³

Time of assessment			
Date of assessment			
Best eye response (E)			
No eye opening	1	1	1
Eye opening in response to pain	2	2	2
Eye opening to speech	3	3	3
Eyes opening spontaneously	4	4	4
Best verbal response (V)			
No verbal response	1	1	1
Incomprehensible sounds	2	2	2
Inappropriate words	3	3	3
Confused	4	4	4
Oriented	5	5	5
Best motor response (M)			
No motor response	1	1	1
Extension to pain	2	2	2
Abnormal flexion to pain	3	3	3
Flexion / Withdrawal to pain	4	4	4
Localizes to pain	5	5	5
Obeys commands	6	6	6
Glasgow Coma score (E + V + M)			

CERVICAL SPINE ASSESSMENT

Does the athlete report that their neck is pain free at rest?	Y	N
If there is NO neck pain at rest , does the athlete have a full range of ACTIVE pain free movement?	Y	N
Is the limb strength and sensation normal?	Y	N

In a patient who is not lucid or fully conscious, a cervical spine injury should be assumed until proven otherwise.

OFFICE OR OFF-FIELD ASSESSMENT

Please note that the neurocognitive assessment should be done in a distraction-free environment with the athlete in a resting state.

STEP 1: ATHLETE BACKGROUND

Sport / team / school: _____

Date / time of injury: _____

Years of education completed: _____

Age: _____

Gender: M / F / Other

Dominant hand: left / neither / right

How many diagnosed concussions has the athlete had in the past?: _____

When was the most recent concussion?: _____

How long was the recovery (time to being cleared to play) from the most recent concussion?: _____ (days)

Has the athlete ever been:

	Yes	No
Hospitalized for a head injury?		
Diagnosed / treated for headache disorder or migraines?		
Diagnosed with a learning disability / dyslexia?		
Diagnosed with ADD / ADHD?		
Diagnosed with depression, anxiety or other psychiatric disorder?		

Current medications? If yes, please list:

Name: _____

DOB: _____

Address: _____

ID number: _____

Examiner: _____

Date: _____

2

STEP 2: SYMPTOM EVALUATION

The athlete should be given the symptom form and asked to read this instruction paragraph out loud then complete the symptom scale. For the baseline assessment, the athlete should rate his/her symptoms based on how he/she typically feels and for the post injury assessment the athlete should rate their symptoms at this point in time.

Please Check: Baseline Post-Injury

Please hand the form to the athlete

	none	mild	moderate	severe			
Headache	0	1	2	3	4	5	6
"Pressure in head"	0	1	2	3	4	5	6
Neck Pain	0	1	2	3	4	5	6
Nausea or vomiting	0	1	2	3	4	5	6
Dizziness	0	1	2	3	4	5	6
Blurred vision	0	1	2	3	4	5	6
Balance problems	0	1	2	3	4	5	6
Sensitivity to light	0	1	2	3	4	5	6
Sensitivity to noise	0	1	2	3	4	5	6
Feeling slowed down	0	1	2	3	4	5	6
Feeling like "in a fog"	0	1	2	3	4	5	6
"Don't feel right"	0	1	2	3	4	5	6
Difficulty concentrating	0	1	2	3	4	5	6
Difficulty remembering	0	1	2	3	4	5	6
Fatigue or low energy	0	1	2	3	4	5	6
Confusion	0	1	2	3	4	5	6
Drowsiness	0	1	2	3	4	5	6
More emotional	0	1	2	3	4	5	6
Irritability	0	1	2	3	4	5	6
Sadness	0	1	2	3	4	5	6
Nervous or Anxious	0	1	2	3	4	5	6
Trouble falling asleep (if applicable)	0	1	2	3	4	5	6

Total number of symptoms: _____ of 22

Symptom severity score: _____ of 132

Do your symptoms get worse with physical activity? Y N

Do your symptoms get worse with mental activity? Y N

If 100% is feeling perfectly normal, what percent of normal do you feel?

If not 100%, why?

Please hand form back to examiner

STEP 3: COGNITIVE SCREENING

Standardised Assessment of Concussion (SAC)⁴

ORIENTATION

What month is it?	0	1
What is the date today?	0	1
What is the day of the week?	0	1
What year is it?	0	1
What time is it right now? (within 1 hour)	0	1
Orientation score	of 5	

IMMEDIATE MEMORY

The Immediate Memory component can be completed using the traditional 5-word per trial list or optionally using 10-words per trial to minimise any ceiling effect. All 3 trials must be administered irrespective of the number correct on the first trial. Administer at the rate of one word per second.

Please choose EITHER the 5 or 10 word list groups and circle the specific word list chosen for this test.

I am going to test your memory. I will read you a list of words and when I am done, repeat back as many words as you can remember, in any order. For Trials 2 & 3: I am going to repeat the same list again. Repeat back as many words as you can remember in any order, even if you said the word before.

List	Alternate 5 word lists					Score (of 5)		
						Trial 1	Trial 2	Trial 3
A	Finger	Penny	Blanket	Lemon	Insect			
B	Candle	Paper	Sugar	Sandwich	Wagon			
C	Baby	Monkey	Perfume	Sunset	Iron			
D	Elbow	Apple	Carpet	Saddle	Bubble			
E	Jacket	Arrow	Pepper	Cotton	Movie			
F	Dollar	Honey	Mirror	Saddle	Anchor			
Immediate Memory Score						of 15		
Time that last trial was completed								

List	Alternate 10 word lists					Score (of 10)		
						Trial 1	Trial 2	Trial 3
G	Finger	Penny	Blanket	Lemon	Insect			
	Candle	Paper	Sugar	Sandwich	Wagon			
H	Baby	Monkey	Perfume	Sunset	Iron			
	Elbow	Apple	Carpet	Saddle	Bubble			
I	Jacket	Arrow	Pepper	Cotton	Movie			
	Dollar	Honey	Mirror	Saddle	Anchor			
Immediate Memory Score						of 30		
Time that last trial was completed								

Name: _____
 DOB: _____
 Address: _____
 ID number: _____
 Examiner: _____
 Date: _____

CONCENTRATION

DIGITS BACKWARDS

Please circle the Digit list chosen (A, B, C, D, E, F). Administer at the rate of one digit per second reading DOWN the selected column.

I am going to read a string of numbers and when I am done, you repeat them back to me in reverse order of how I read them to you. For example, if I say 7-1-9, you would say 9-1-7.

Concentration Number Lists (circle one)					
List A	List B	List C			
4-9-3	5-2-6	1-4-2	Y	N	0
6-2-9	4-1-5	6-5-8	Y	N	1
3-8-1-4	1-7-9-5	6-8-3-1	Y	N	0
3-2-7-9	4-9-6-8	3-4-8-1	Y	N	1
6-2-9-7-1	4-8-5-2-7	4-9-1-5-3	Y	N	0
1-5-2-8-6	6-1-8-4-3	6-8-2-5-1	Y	N	1
7-1-8-4-6-2	8-3-1-9-6-4	3-7-6-5-1-9	Y	N	0
5-3-9-1-4-8	7-2-4-8-5-6	9-2-6-5-1-4	Y	N	1
List D	List E	List F			
7-8-2	3-8-2	2-7-1	Y	N	0
9-2-6	5-1-8	4-7-9	Y	N	1
4-1-8-3	2-7-9-3	1-6-8-3	Y	N	0
9-7-2-3	2-1-6-9	3-9-2-4	Y	N	1
1-7-9-2-6	4-1-8-6-9	2-4-7-5-8	Y	N	0
4-1-7-5-2	9-4-1-7-5	8-3-9-6-4	Y	N	1
2-6-4-8-1-7	6-9-7-3-8-2	5-8-6-2-4-9	Y	N	0
8-4-1-9-3-5	4-2-7-9-3-8	3-1-7-8-2-6	Y	N	1
Digits Score:					of 4

MONTHS IN REVERSE ORDER

Now tell me the months of the year in reverse order. Start with the last month and go backward. So you'll say December, November. Go ahead.

Dec - Nov - Oct - Sept - Aug - Jul - Jun - May - Apr - Mar - Feb - Jan	0	1
Months Score	of 1	
Concentration Total Score (Digits + Months)	of 5	

4

STEP 4: NEUROLOGICAL SCREEN

See the instruction sheet (page 7) for details of test administration and scoring of the tests.

Can the patient read aloud (e.g. symptom checklist) and follow instructions without difficulty?	Y	N
Does the patient have a full range of pain-free PASSIVE cervical spine movement?	Y	N
Without moving their head or neck, can the patient look side-to-side and up-and-down without double vision?	Y	N
Can the patient perform the finger nose coordination test normally?	Y	N
Can the patient perform tandem gait normally?	Y	N

BALANCE EXAMINATION

Modified Balance Error Scoring System (mBESS) testing⁵

Which foot was tested (i.e. which is the non-dominant foot) Left Right

Testing surface (hard floor, field, etc.) _____

Footwear (shoes, barefoot, braces, tape, etc.) _____

Condition	Errors
Double leg stance	_____ of 10
Single leg stance (non-dominant foot)	_____ of 10
Tandem stance (non-dominant foot at the back)	_____ of 10
Total Errors	_____ of 30

Name: _____

DOB: _____

Address: _____

ID number: _____

Examiner: _____

Date: _____

5

STEP 5: DELAYED RECALL:

The delayed recall should be performed after 5 minutes have elapsed since the end of the Immediate Recall section. Score 1 pt. for each correct response.

Do you remember that list of words I read a few times earlier? Tell me as many words from the list as you can remember in any order.

Time Started

Please record each word correctly recalled. Total score equals number of words recalled.

Total number of words recalled accurately: of 5 or of 10

6

STEP 6: DECISION

Domain	Date & time of assessment:		
	_____	_____	_____
Symptom number (of 22)	_____	_____	_____
Symptom severity score (of 132)	_____	_____	_____
Orientation (of 5)	_____	_____	_____
Immediate memory	_____ of 15 _____ of 30	_____ of 15 _____ of 30	_____ of 15 _____ of 30
Concentration (of 5)	_____	_____	_____
Neuro exam	Normal Abnormal	Normal Abnormal	Normal Abnormal
Balance errors (of 30)	_____	_____	_____
Delayed Recall	_____ of 5 _____ of 10	_____ of 5 _____ of 10	_____ of 5 _____ of 10

Date and time of injury: _____

If the athlete is known to you prior to their injury, are they different from their usual self?

Yes No Unsure Not Applicable

(If different, describe why in the clinical notes section)

Concussion Diagnosed?

Yes No Unsure Not Applicable

If re-testing, has the athlete improved?

Yes No Unsure Not Applicable

I am a physician or licensed healthcare professional and I have personally administered or supervised the administration of this SCAT5.

Signature: _____

Name: _____

Title: _____

Registration number (if applicable): _____

Date: _____

SCORING ON THE SCAT5 SHOULD NOT BE USED AS A STAND-ALONE METHOD TO DIAGNOSE CONCUSSION, MEASURE RECOVERY OR MAKE DECISIONS ABOUT AN ATHLETE'S READINESS TO RETURN TO COMPETITION AFTER CONCUSSION.

CLINICAL NOTES:

Name: _____
 DOB: _____
 Address: _____
 ID number: _____
 Examiner: _____
 Date: _____



CONCUSSION INJURY ADVICE

(To be given to the person monitoring the concussed athlete)

This patient has received an injury to the head. A careful medical examination has been carried out and no sign of any serious complications has been found. Recovery time is variable across individuals and the patient will need monitoring for a further period by a responsible adult. Your treating physician will provide guidance as to this timeframe.

If you notice any change in behaviour, vomiting, worsening headache, double vision or excessive drowsiness, please telephone your doctor or the nearest hospital emergency department immediately.

Other important points:

Initial rest: Limit physical activity to routine daily activities (avoid exercise, training, sports) and limit activities such as school, work, and screen time to a level that does not worsen symptoms.

- 1) Avoid alcohol
- 2) Avoid prescription or non-prescription drugs without medical supervision. Specifically:
 - a) Avoid sleeping tablets
 - b) Do not use aspirin, anti-inflammatory medication or stronger pain medications such as narcotics
- 3) Do not drive until cleared by a healthcare professional.
- 4) Return to play/sport requires clearance by a healthcare professional.

Clinic phone number: _____

Patient's name: _____

Date / time of injury: _____

Date / time of medical review: _____

Healthcare Provider: _____

© Concussion in Sport Group 2017

Contact details or stamp

INSTRUCTIONS

Words in *Italics* throughout the SCAT5 are the instructions given to the athlete by the clinician

Symptom Scale

The time frame for symptoms should be based on the type of test being administered. At baseline it is advantageous to assess how an athlete "typically" feels whereas during the acute/post-acute stage it is best to ask how the athlete feels at the time of testing.

The symptom scale should be completed by the athlete, not by the examiner. In situations where the symptom scale is being completed after exercise, it should be done in a resting state, generally by approximating his/her resting heart rate.

For total number of symptoms, maximum possible is 22 except immediately post injury, if sleep item is omitted, which then creates a maximum of 21.

For Symptom severity score, add all scores in table, maximum possible is 22 x 6 = 132, except immediately post injury if sleep item is omitted, which then creates a maximum of 21x6=126.

Immediate Memory

The Immediate Memory component can be completed using the traditional 5-word per trial list or, optionally, using 10-words per trial. The literature suggests that the Immediate Memory has a notable ceiling effect when a 5-word list is used. In settings where this ceiling is prominent, the examiner may wish to make the task more difficult by incorporating two 5-word groups for a total of 10 words per trial. In this case, the maximum score per trial is 10 with a total trial maximum of 30.

Choose one of the word lists (either 5 or 10). Then perform 3 trials of immediate memory using this list.

Complete all 3 trials regardless of score on previous trials.

"I am going to test your memory. I will read you a list of words and when I am done, repeat back as many words as you can remember, in any order." The words must be read at a rate of one word per second.

Trials 2 & 3 MUST be completed regardless of score on trial 1 & 2.

Trials 2 & 3:

"I am going to repeat the same list again. Repeat back as many words as you can remember in any order, even if you said the word before."

Score 1 pt. for each correct response. Total score equals sum across all 3 trials. Do NOT inform the athlete that delayed recall will be tested.

Concentration

Digits backward

Choose one column of digits from lists A, B, C, D, E or F and administer those digits as follows:

Say: *"I am going to read a string of numbers and when I am done, you repeat them back to me in reverse order of how I read them to you. For example, if I say 7-1-9, you would say 9-1-7."*

Begin with first 3 digit string.

If correct, circle "Y" for correct and go to next string length. If incorrect, circle "N" for the first string length and read trial 2 in the same string length. One point possible for each string length. Stop after incorrect on both trials (2 N's) in a string length. The digits should be read at the rate of one per second.

Months in reverse order

"Now tell me the months of the year in reverse order. Start with the last month and go backward. So you'll say December, November ... Go ahead"

1 pt. for entire sequence correct

Delayed Recall

The delayed recall should be performed after 5 minutes have elapsed since the end of the Immediate Recall section.

"Do you remember that list of words I read a few times earlier? Tell me as many words from the list as you can remember in any order."

Score 1 pt. for each correct response

Modified Balance Error Scoring System (mBESS)⁵ testing

This balance testing is based on a modified version of the Balance Error Scoring System (BESS)⁵. A timing device is required for this testing.

Each of 20-second trial/stance is scored by counting the number of errors. The examiner will begin counting errors only after the athlete has assumed the proper start position. The modified BESS is calculated by adding one error point for each error during the three 20-second tests. The maximum number of errors for any single condition is 10. If the athlete commits multiple errors simultaneously, only

one error is recorded but the athlete should quickly return to the testing position, and counting should resume once the athlete is set. Athletes that are unable to maintain the testing procedure for a minimum of five seconds at the start are assigned the highest possible score, ten, for that testing condition.

OPTION: For further assessment, the same 3 stances can be performed on a surface of medium density foam (e.g., approximately 50cm x 40cm x 6cm).

Balance testing – types of errors

- | | | |
|---------------------------------|---|---|
| 1. Hands lifted off iliac crest | 3. Step, stumble, or fall | 5. Lifting forefoot or heel |
| 2. Opening eyes | 4. Moving hip into > 30 degrees abduction | 6. Remaining out of test position > 5 sec |

"I am now going to test your balance. Please take your shoes off (if applicable), roll up your pant legs above ankle (if applicable), and remove any ankle taping (if applicable). This test will consist of three twenty second tests with different stances."

(a) Double leg stance:

"The first stance is standing with your feet together with your hands on your hips and with your eyes closed. You should try to maintain stability in that position for 20 seconds. I will be counting the number of times you move out of this position. I will start timing when you are set and have closed your eyes."

(b) Single leg stance:

"If you were to kick a ball, which foot would you use? [This will be the dominant foot] Now stand on your non-dominant foot. The dominant leg should be held in approximately 30 degrees of hip flexion and 45 degrees of knee flexion. Again, you should try to maintain stability for 20 seconds with your hands on your hips and your eyes closed. I will be counting the number of times you move out of this position. If you stumble out of this position, open your eyes and return to the start position and continue balancing. I will start timing when you are set and have closed your eyes."

(c) Tandem stance:

"Now stand heel-to-toe with your non-dominant foot in back. Your weight should be evenly distributed across both feet. Again, you should try to maintain stability for 20 seconds with your hands on your hips and your eyes closed. I will be counting the number of times you move out of this position. If you stumble out of this position, open your eyes and return to the start position and continue balancing. I will start timing when you are set and have closed your eyes."

Tandem Gait

Participants are instructed to stand with their feet together behind a starting line (the test is best done with footwear removed). Then, they walk in a forward direction as quickly and as accurately as possible along a 38mm wide (sports tape), 3 metre line with an alternate foot heel-to-toe gait ensuring that they approximate their heel and toe on each step. Once they cross the end of the 3m line, they turn 180 degrees and return to the starting point using the same gait. Athletes fail the test if they step off the line, have a separation between their heel and toe, or if they touch or grab the examiner or an object.

Finger to Nose

"I am going to test your coordination now. Please sit comfortably on the chair with your eyes open and your arm (either right or left) outstretched (shoulder flexed to 90 degrees and elbow and fingers extended), pointing in front of you. When I give a start signal, I would like you to perform five successive finger to nose repetitions using your index finger to touch the tip of the nose, and then return to the starting position, as quickly and as accurately as possible."

References

1. McCrory et al. Consensus Statement On Concussion In Sport – The 5th International Conference On Concussion In Sport Held In Berlin, October 2016. British Journal of Sports Medicine 2017 (available at www.bjsm.bmj.com)
2. Maddocks, DL; Dicker, GD; Saling, MM. The assessment of orientation following concussion in athletes. Clinical Journal of Sport Medicine 1995; 5: 32-33
3. Jennett, B., Bond, M. Assessment of outcome after severe brain damage: a practical scale. Lancet 1975; i: 480-484
4. McCrea M. Standardized mental status testing of acute concussion. Clinical Journal of Sport Medicine. 2001; 11: 176-181
5. Guskiewicz KM. Assessment of postural stability following sport-related concussion. Current Sports Medicine Reports. 2003; 2: 24-30

CONCUSSION INFORMATION

Any athlete suspected of having a concussion should be removed from play and seek medical evaluation.

Signs to watch for

Problems could arise over the first 24-48 hours. The athlete should not be left alone and must go to a hospital at once if they experience:

- Worsening headache
- Drowsiness or inability to be awakened
- Inability to recognize people or places
- Repeated vomiting
- Unusual behaviour or confusion or irritable
- Seizures (arms and legs jerk uncontrollably)
- Weakness or numbness in arms or legs
- Unsteadiness on their feet.
- Slurred speech

Consult your physician or licensed healthcare professional after a suspected concussion. Remember, it is better to be safe.

Rest & Rehabilitation

After a concussion, the athlete should have physical rest and relative cognitive rest for a few days to allow their symptoms to improve. In most cases, after no more than a few days of rest, the athlete should gradually increase their daily activity level as long as their symptoms do not worsen. Once the athlete is able to complete their usual daily activities without concussion-related symptoms, the second step of the return to play/sport progression can be started. The athlete should not return to play/sport until their concussion-related symptoms have resolved and the athlete has successfully returned to full school/learning activities.

When returning to play/sport, the athlete should follow a stepwise, **medically managed exercise progression, with increasing amounts of exercise.** For example:

Graduated Return to Sport Strategy

Exercise step	Functional exercise at each step	Goal of each step
1. Symptom-limited activity	Daily activities that do not provoke symptoms.	Gradual reintroduction of work/school activities.
2. Light aerobic exercise	Walking or stationary cycling at slow to medium pace. No resistance training.	Increase heart rate.
3. Sport-specific exercise	Running or skating drills. No head impact activities.	Add movement.
4. Non-contact training drills	Harder training drills, e.g., passing drills. May start progressive resistance training.	Exercise, coordination, and increased thinking.
5. Full contact practice	Following medical clearance, participate in normal training activities.	Restore confidence and assess functional skills by coaching staff.
6. Return to play/sport	Normal game play.	

In this example, it would be typical to have 24 hours (or longer) for each step of the progression. If any symptoms worsen while exercising, the athlete should go back to the previous step. Resistance training should be added only in the later stages (Stage 3 or 4 at the earliest).

Written clearance should be provided by a healthcare professional before return to play/sport as directed by local laws and regulations.

Graduated Return to School Strategy

Concussion may affect the ability to learn at school. The athlete may need to miss a few days of school after a concussion. When going back to school, some athletes may need to go back gradually and may need to have some changes made to their schedule so that concussion symptoms do not get worse. If a particular activity makes symptoms worse, then the athlete should stop that activity and rest until symptoms get better. To make sure that the athlete can get back to school without problems, it is important that the healthcare provider, parents, caregivers and teachers talk to each other so that everyone knows what the plan is for the athlete to go back to school.

Note: If mental activity does not cause any symptoms, the athlete may be able to skip step 2 and return to school part-time before doing school activities at home first.

Mental Activity	Activity at each step	Goal of each step
1. Daily activities that do not give the athlete symptoms	Typical activities that the athlete does during the day as long as they do not increase symptoms (e.g. reading, texting, screen time). Start with 5-15 minutes at a time and gradually build up.	Gradual return to typical activities.
2. School activities	Homework, reading or other cognitive activities outside of the classroom.	Increase tolerance to cognitive work.
3. Return to school part-time	Gradual introduction of school-work. May need to start with a partial school day or with increased breaks during the day.	Increase academic activities.
4. Return to school full-time	Gradually progress school activities until a full day can be tolerated.	Return to full academic activities and catch up on missed work.

If the athlete continues to have symptoms with mental activity, some other accommodations that can help with return to school may include:

- Starting school later, only going for half days, or going only to certain classes
- More time to finish assignments/tests
- Quiet room to finish assignments/tests
- Not going to noisy areas like the cafeteria, assembly halls, sporting events, music class, shop class, etc.
- Taking lots of breaks during class, homework, tests
- No more than one exam/day
- Shorter assignments
- Repetition/memory cues
- Use of a student helper/tutor
- Reassurance from teachers that the child will be supported while getting better

The athlete should not go back to sports until they are back to school/learning, without symptoms getting significantly worse and no longer needing any changes to their schedule.

Post-Concussion Symptom Scale

Athlete Name: _____ DOI: _____

Symptom	Date: _____ Time: _____	Date: _____ Time: _____	Date: _____ Time: _____	Date: _____ Time: _____	Date: _____ Time: _____	Date: _____ Time: _____	Date: _____ Time: _____
	Headache						
"Pressure in head"							
Neck pain							
Balance problems or dizziness							
Nausea or vomiting							
Vision Problems							
Hearing problems / ringing							
"Don't feel right"							
Feeling "dinged" or "dazed"							
Confusion							
Feeling slowed down							
Feeling like "In a fog"							
Drowsiness							
Fatigue or low energy							
More emotional than usual							
Irritability							
Difficulty concentrating							
Difficulty remembering							
Sadness							
Nervous or anxious							
Trouble falling asleep							
Sleeping more than usual							
Sensitivity to light							
Sensitivity to noise							
Other _____							

The Post-Concussion Symptom Scale should be used for the initial evaluation on the SCAT evaluation and for each subsequent follow-up assessment until all signs/symptoms have cleared at rest and during physical exertion. The athletic trainer will ask the athlete to grade or score the severity of the symptom on a scale of 0-6, where 0 = not present, 1= mild, 3 = moderate, and 6 = most severe.

Post-Concussion Symptom Scale

Athlete Name: _____ DOI: _____

Symptom	Date: _____ Time: _____	Date: _____ Time: _____	Date: _____ Time: _____	Date: _____ Time: _____	Date: _____ Time: _____	Date: _____ Time: _____	Date: _____ Time: _____
	Headache						
"Pressure in head"							
Neck pain							
Balance problems or dizziness							
Nausea or vomiting							
Vision Problems							
Hearing problems / ringing							
"Don't feel right"							
Feeling "dinged" or "dazed"							
Confusion							
Feeling slowed down							
Feeling like "In a fog"							
Drowsiness							
Fatigue or low energy							
More emotional than usual							
Irritability							
Difficulty concentrating							
Difficulty remembering							
Sadness							
Nervous or anxious							
Trouble falling asleep							
Sleeping more than usual							
Sensitivity to light							
Sensitivity to noise							
Other _____							

The Post-Concussion Symptom Scale should be used for the initial evaluation on the SCAT evaluation and for each subsequent follow-up assessment until all signs/symptoms have cleared at rest and during physical exertion. The athletic trainer will ask the athlete to grade or score the severity of the symptom on a scale of 0-6, where 0 = not present, 1= mild, 3 = moderate, and 6 = most severe.